
32-year-old male
history of trauma or operations
edical history : irrelevant
al dendritic ulcer(linear central corneal ulcer)
هيربس



e main symptom is severe pain. F
e causative agent is fungal . F
rneal perforation is a rare complication. T
pical steroids are contraindicated . T
pical broad-spectrum antibiotics is the main line of treatment F
currence after cure of the condition is rare .F
ypopyon is commonly associated with this condition F

A 48-year-old female presented with sudden red painful eye and drop of vision جلوكوما Acute angle Closure glauc



The condition can be managed on out-patient basis. F
Topical steroids should be initiated quickly .F
The expected refraction is high hypermetropia.T
Surgical management is usually indicated .T
Topical miotics are indicated .T
The fellow eye is not at risk of developing similar condition.F
Topical mydriatics are contraindicated .T

A 14-year old boy presented with **sticky** discharge
gluing the lashes together
no photophobia.
vision is 6/6 OU and pupils are normal and reactive
باكتريل كونجكتفيتز **Bact conjunctivitis**



arding this condition :

he causative agent is mostly viral .
atching the eye is essential.F
leasuring IOP is not necessary . T
ntibiotic eye drops is the main line of treatement T
ledical systemic work up is indicated.F

18-years old boy

Bact conjunctivitis



Regarding this condition

1. It is related to poor hygiene T
2. Patient can be assured it is not contagiousF
3. Topical steroids are the mainstay of management F
4. Corneal ulcer is a potential complication T
5. *Systemic antibiotics are routinely prescribed F*

A 20-year old man in military camp
Presented with watery eye and Mild photophobia.
vision is 6/18 OU and pupils are normal and reactive
Viral conjunctivitis



rding this condition :

ne causative agent is mostly viral .
tching the eye is essential.F
easuring IOP is essential . F
ntibiotic eye drops is the main line of treatment F
ne condition is usually sporadic F
ne condition is usually self-limiting.T

This case presented with unilateral red aching eye

Phlyctenular conjunctivitis



xternal allergy sensitivity to endogenous agent

rding this condition :

s common in old age .F

e underlying pathogenesis is allergy to sunlight.F

pical steroids are the main line of treatment . T

e corneal can be involved in this condition .T بس الكتاب يقول انها نادره تحصل


e condition is usually self-limiting. F

stemic workup is needed T

A 56-years-old woman presented with pain ,

photophobia and drop of vision .

Bact corneal ulcer or Fungal ulcer



The predisposing factors :

1. Corneal anesthesi T
2. Kerato- conjunctivitis sic a .T
3. Lagophthalmos with expos e T.
4. Long term topical anti – glaucoma eye drops.F
5. Ptosis. F

A 32-year-old male presented with red , pain
eye and drop of vision .
He gave a history of oral and genital ulcers

Behcet

Ant. uveitis



Regarding the management of this condition

1. Topical steroids
2. Systemic carbonic anhydrase inhibitors.F
3. Topical antibiotic ointment F
4. Topical miotics
5. Topical cycloplegics.T
6. Rheumatology consultation . T

34-years-old man
presented with acute painful red
Ant. uveitis



1- Topical miotics are indicated in treating this patient.F
2- Topical steroids are contraindicated .F
3- The AC shows hyphema .F
4- Systemic associations should be considered . T
5- Plain X-ray to the sacro-iliac joint and pelvis is indicated .T


A 19-year-old female
History of contact lens wear
Acanthameba keratitis



1- The main symptom is severe pain out of proportion of signs.T
2- Bad hygiene has no role in the etiology of this condition .F
3- The condition is usually associated with radial peri neuritis T.
4- Propamidine Isethianate is the main line of

A 32-year-old man with high myopia

Retina detachment



l complaint of the patient is flashes of light and floaters.T


diagnosis is tractional retinal detachment .F

ly history may be one of the risk factors .T

retinal break can be treated with argon laser photocoagulatio .F


palna vitrectomy has no role in management of this case. F

A 55-year-old diabetic patient
IOP : 35 mmHg
Rubeosis iridis




1- Fundus examination may reveal non-proliferative diabetic retinopathy .F
2- Peripheral anterior synechiae is the cause of developing glaucoma.T
3- Conventional glaucoma surgery is likely to fail in this condition.T
4- Pan retinal photo-coagulation is indicated in this case T
5- The condition is secondary to retinal ischemia

A 32-year-old man with a history of severe head injury , was admitted to the hospital for status post concussion .



Subconjunctival hge


1- Reassurance and decongestant eye drops are prescribed F.
2- Neurologic consultation and neuro imaging are require T.
3- Fundus examination is not required in this case F
4- Topical antibiotics are the primary line of treatment .F
5- Plain X ray skull is essential in evaluating this cas .T



This ocular disease is most commonly associated with the following systemic disease nodular Episcleritis


1. Hyper thyroisism
2. Behcet’s disease
3. Sarcoidosis
4. Rheumatoid arthritisT
5. Tuberculosis .

A 65-year-old male
Presented with epiphora
and chronic irritation
Entropion



1- Topical lubricant eye drops can alleviate the condition .
2- Corneal staining with Fluorescein is necessary T
3- Scarring of the palpebral conjunctiva can be the causeT.
4- Exophthalmos is a predisposing factor .F
5- Surgical management is quickly recommended .


A 14-year-old boy
VKCالـرمد الـربيعي



1- The underlying pathogenesis is allergy to exogenous agents .T
2- Keratoconus may be associated .T
3- Muco purulent discharge is usually present.F Roping discharge
4- Long-term topical steroid eye drops are the first line of treatment .f In severe cases
5- Itching is usually not prominent .F

38-year-old man presented with
sfigurement and chronic visual lo

erygium



1- Long-term exposure to UV light and dust is a predisposing facto T

2- Medical treatment is indicated in this ase.F

3- Recurrence is rare after surgical e ision F.

4- The condition is more common in m es .T

5- The condition is usually statio ary .F

A 52-old-female patient
presented with blurring of vision

diabetic retinopathy



1- Systemic work-up includes HbA1c

2- OCT and Fluorescein angiography is useful in diagnosis T

3- Laser photocoagulation is indicated .T


4- The duration of systemic illness is an important risk factor .T

5- Reassurance and routine follow up are recommended.F

A 60-year-old male

Nasal shift- cupping of visual disc

Glaucoma



1- Measurement of IOP is not essential in this case.

2- Blurring of vision is an early symptom in this condition.

3- Visual field examination is not important in this conditio F.


4- Optical Coherence Tomography is one of the investigative tools to follow up this case.

5- Trabeculectomy is the standard surgery in this conditio F .


لازم ابداء الاول بالادويه

A 70-year-old male complains of drop of vision in his
right eye since 1 week. He noticed that the straight lines
seemed broken **Metamorphopsia**

ARMD




1- The underlying pathogenesis is infectious .F
2- Ultrasonography is important to confirm the diagnosis .F
3- Argon laser photocoagulation has no role in management T.
4- Intra vitreal Anti – VEGF is indicated in this case T
5- Amsler grid is useful in diagnosis and follow up T
سبه للارجون ليزر هو قال فى المحاضره انه له دور كبير جدا ؟؟؟



A 36-year old male with type I diabetes mellitus

proliferative retinopathy



The duration of diabetes is an important risk factor.T


Strict diabetic control has no role in prevention of this condition.F

Argon laser photo coagulation could have prevented this condition.T

The patient is at risk of vitreous hemorrhage . T

Retinal ischemia is the cause of this picture.T

47-years -old female
Type II diabetes



DIABETIC MACULOPATHY

Regarding this condition

1. The diagnosis is proliferative diabetic retinopathy.F
2. Initial management is pan retinal photocoagulationF
3. Strict diabetic control alone can cure this condition F
4. The duration of diabetes is an important risk factor T
5. The underlying pathology in this case is capillary occlusion .F

ب علاج هبدأ ANTI- VEGF

- والسبب CAPILLAY LEAKAGE

3- years - old female
ype I (IDDM) diabetes for 15 years
bA1c 9.5%

RE-RETINAL VITEROUS HEMORRAGHE

nagement of this condition includes :

anti-platelet drugsF
Ultrasonography T
Fundus fluorescein angiographyT
Laser photocoagulationT
Diabetic control and follow upT

بالنسبه لرقم 2 مكتوبه فى عين شمس هستخ B-SCAN



A 20-year old female with acute drop of vision to

6/60 OD, lost color vision. MS



1- Visual evoked potential is useful to reach the diagnosis .T

2- MRI of the brain is important .T

3- The condition is usually painless .F

4- Pupillary reflexes are essential during examination .T

5- Oral steroid therapy is the mainstay of management .F



بيتم فى 5 دقائق والطبيعى بيكون من 30-10 مم

Regarding this clinical test the following is true :**Shcimmer test**

1. It is evaluation of corneal sensitivityF
2. It is done in dark room F
3. It is evaluated after waiting for 15 minutesF
4. The values are expressed in millimeters T
5. It is done with and without topical

48-year-old man presented with drop of vision , loss of weight and low-grade fever


- Leukemia **roth spot hge**



The systemic workup of this patient :

1. Glyclosylated hemoglobin (Hb A1c) F
2. Blood pressure measurement F
3. Endocrinological consultation F
4. Complete Blood count and sternal puncture .T
5. Immunological consultation .f

This is the fundus picture of 72 - years -old patient , presented with sudden severe drop of vision to 1/60 He complains of general fatigue and fever.




GIANT CELL ARTHERITIS

Regarding this condition :

1. Lumber puncture and CSF study might help to reach the diagnosis F.
2. Intra venous steroid therapy is sight - saving . T
3. ESR is usually within normal . F
4. *The visual field showed concentric contraction .F*

Field defect present; commonly ALTITUDINAL

5. Carotid Doppler study is important to confirm the



Optic nerve head of a 19-years-old girl . She presented to the outpatient clinic for routine check up . Pupillary light reflexes were normal in both eyes and no RAPD **Papilledema**

The next step in management include :

1. Fundus Fluorescein Angiography and macular OCT. F
2. Visual field test . T
3. Observation and repeated fundus examination .T
4. Systemic steroid therapy. F

Optic nerve head of a 54-year-old female
Attending OP clinic for routine check – up
Glaucoma



For management of this case :

1. Visual field test is essential T
2. Family history is irrelevant in this condition F
3. Reassurance and follow up of the case .F
4. Refer to neurologist for CSF pressure evaluation F.
5. MRI brain is needed to reach the diagnosis .F

- Disc changes in Glaucoma:
  - Increased cup / disc ratio.
  - Vertical elongation of cup
  - Visibility of pores of lamina cribrosa
  - Nasal shift of disc vessels
  - Bayonetting , notching
  - Splinter hemorrhage
  - Peripapillary pigmentation
  - Cup asymmetry

A 27-years-old female presented with sudden drop of vision in the right eye to 6/60 . The left eye was 6/6 vision and ophthalmologically -free There was deep seated retro bulbar pain with eye movements and RAPD in the right eye.


Retrobulbar neuritis



The next step in management include :

1. Serial lumbar puncture F
2. MRI and neurological consultatio . T
3. Urgent intra venous pulsed steroid the py .T
4. Orbital decompression. F
5. Systemic diuretics as Lasix. F

Slit-lamp image and plain chest
X-ray of a 58-year-old male
patient, complaining of low grade pain and gradual
drop of vision in his left eye
Sarcoidosis
.





Regarding this case :

1. Angiotensin converting enzyme (ACE) is important to confirm the diagnosis t
2. It is due to chronic granulomatous systemic disease T
3. Dermatologic affection is rare in this condition .F
4. Systemic broad – spectrum antibiotics should be started immediately .F
5. It is more in common in blacks .T

This 32-years-old man presented with chronic redness , lacrimation   sjs




The diagnosis can be reached by :

1. Intra ocular pressure (IOP) measurements. F
2. Fundus examination and fluorescein angiography.F
3. History of recent drug intake.T
4. Corneal endothelial cell count . F
5. Conjunctival smear / biopsy . T

A 65-year-old man presented with sudden loss

Of upper visual fields 3 days ago



Branch retinal artery occlusion

Management of this condition includes :


1. anti-coagulants .T
2. Ocualr massage T
3. Ultrasonography F
4. Fundus fluorescein angiographyT
5. Cardiological consultationT

A 24-years old man presented with drop of vision

in the right eye .

Examination revealed tall person with long

arms and long slender fingers . **Marfan**



1- The patient may complain of binocular diplopia .F MONOCULAR NOT BINOCUL

2- The whole zonular attachment of the lens is lost.F


3- Cardiological consultation is important .

4- Glass prescription may relief the patient’s complaint.

5- The mode of inherence is autosomal dominant T

A 58-year-old female presented with sudden drop of vision 7 days ago

crvo



Regarding this condition :

1. Initial diagnosis is proliferaive diabetic retinopathy F .
2. Laser therapy is indicated immediately F monthly injection of anti-VEGF for 6-8 months
3. Measuring IOP is not necessary .T
4. Fundus fluorescein angiography is indicated T
5. Medical systemic work up is indicated

A 54-year old woman
Medial history rheumatoid artheritis
Presented with irritation , FB sensation , lacrimation .
vision is 6/18 and pupils are normal and reactive

Dry eye




garding this condition :

All false

Ocular ultrasonography is important to confirm the diagnosis F   .
Patching the eye is essentialF
Measuring IOP is essential . F
Antibiotic eye drops is the main line of treatment
The condition is usually self-limiting. F

A 31-year old man
Medial history low back ache and kyphosis
Presented with redness , pain
vision is 6/24 **Iridocyclitis 2ry to Ankylosing Spond**



ding this condition :

ocular condition is usually recurrent T
condition is usually associated with HLA B5
suring IOP is essential . **T**
Bec it may becompliacted with glaucoma?
biotic eye drops is the main line of treatment
diological consultation is essential . F

A 45-year old female complains change of
Appearance over the past few years
Serum TSH is decreased below normal level.



1- MRI shows bilateral enlargement of the media and inferior re .T
2- Cigarette smoking has no relation to this condition .
3- Ocular manifestations clears when the thyroid functions are normal ed F.
4- Optic nerve compression may be the cause of severe visual l s. T
5- Systemic steroid may be sight-saving in some cases .

A 5-year-old boy leukocoria



The following diseases can be the cause of this appearance

1. Keratoconu .F
2. Congenital cata ct T
3. Sub luxtated l s F
4. Retinoblasto a T
5. Coats' diseas . T

A 6-year old boy with fair hair and skin

Homocysteinuria




the child needs surgery , general anesthesia is s


Mental retardation is sometimes associated .T

ens zonules are usually stretched. F

erum homocystein is important investigation



Visual field of a 65-year-old man with right field defect non artritc



in anterior segment examination , the only positive sign will be RAPD

1- The patient has normal pupillary reaction in the right eye .F

2- ESR is usually normal .T


3- Fundus examination will show pale disc edema on the right eye T.

other eye has a CROWDED small di


4- The patient is at risk of similar condition in the left eye later. T

5- This visual field defect is called homonymous hemi anopia .F

Causes of this fundus appearance include :



1- Central retinal artery occlusion.T
2- Blunt Trauma.T
3- Central Retinal Vein Occlusion .F
3- Tay-Sachs disease. T
4- Age related macular Degeneration.F



6- Patients with this error of refraction complains of defective near vision .

7- This condition is corrected with concave lenses.


8- Retinal detachment can occur in high degrees.

9- The condition is associated with long axial length .

10- High degrees is associated with apparent convergent squint. T

Apparent esotropia
True exotropia

Visual field of a 65-year-old man
Medial history : hypertension
Heavy smoker and dyslipedemia



Regarding this visual field :

41- This visual field defect is termed homonymous hemi anopia

42- The neurological lesion is expected to be in the left parietal lobe

43- Pupillary reaction to light is absent in both eyes

44- Visual acuity is slightly affected .

45- Plain skull X-ray is needed to reach the diagnosis .

This 17 -years-old boy presented with history of blunt trauma



Regarding this condition :

61- Long-term measurements of Intra ocular pressure (IOP) is needed . T


62- Monocular dilopia is expected .

63- Optical correction by glasses can solve the problem.

64- Fundus examination is required in this case.

65- Corneal topography and pachymetry are helpful for the diagnosis .


A 34-years-old woman presented with acute painful red eye and marked drop of vision .



The management of this case includes :

71- Miotics and hyperosmotics.
72- Local atropine and antibiotics.
73- Local atropine and steroids.
74- Carbonic anhydrase inhibitors.
75- Management on outpatient basis.

A 60-year-old had **sudden** darkening of vision in his left eye.



Regarding this condition

76-The underlying cause is vasculitis.

Occlusion of a retinal artery is usually thrombotic in old age and embolic/vasculitic in young age

77- Ocular massage and intravenous acetazolamide are quickly indicated

78- Carotid Doppler is recommended

79- Visual acuity is expected to be between 6/18 and 6/36

80- Pupillary reaction to light is within normal .

A 17-year-old male



81- The underlying pathogenesis is acute suppurative inflammation of the lash follicle.

82 - Blephartitis is commonly associated .

83- The lesion commonly points to the skin side. conjunctival side

84- Uncorrected error of refraction may be the underlying cause of repeated similar condition .

85- Surgical management by horizontal skin incision .

A 32-old-female patient
complaining of progressive diminution of vision , the patient is looking downwards. Moon sen sign keratocouns



86- The expected refraction is irregular myopic astigmatism .
87- Spectacle correction can improve the vision .
88- Optical biometry is helpful to confirm the diagnosis.
89- Sudden drop of vision can occur in advanced cases .

A 52-old-female patient
presented with blurring of
vision



91- Systemic work-up includes HbA1c .

92- OCT and Fluorescein angiography is useful in diagnosis


93- Laser photocoagulation is indicated .

94- The duration of systemic illness is an important risk factor .

95- Intra vitreal Anti VEGF has no role in

Indicate the use of this test in Ophthalmic Practice:

A. Measure the Corneal Curvature.
B. <u>Measure IOP in mmHg by indentation.</u>
C. Measure Corneal thickness.
D. Measure IOP by Aplantion.
E. Measure the degree of exophthalmos.




Optic Nerve Photograph showing:

A. Normal Optic Nerve Head.
B. Papilledema.
C. Papillitis.
D. <u>Glaucoma Disc changes with Atrophy.</u>
E. Optic Atrophy.



Select the best diagnosis to this Cataract case.

A. Congenital Cataract.
B. Immature Senile Cataract.
C. Traumatic Cataract.
D. Hyper-mature senile Cataract.
E. Mature senile Cataract.



Fundus Photograph of a case chronic visual loss?

A. Chronic Simple Adult Open Angle Glaucoma.
B. Age related macular degeneration, dry type.
C. Age related macular degeneration, wet type.
D. Typical case of Retinitis Pigmentosa.
E. Retinitis Pigmentosa sin Pimento.




Aly has 9 months, with left convergence started after one month of age without limitations of Ocular movement.

a) Follow up is required.
b) Refraction is required using Atropine and refractive error correction.
c) Orthopedic training is required.




23-years old male
History of severe head injury



The next step in management :

a- Topical medical treatment and warm compresses.
<u>b- Neurological consultation and neuro imaging.</u>
c- Systemic steroid therapy.

54-years old female
Drop of vision in left eye




Underlying systemic diseases include the following EXCEPT :

a- Diabetes Mellitus
b- Rheumatoid arthritis
c- Hypertension

73 years old female
Presented with metamorphopsia
and difficult reading


Wet AMD
CNV



Management includes :

<u>a- Repeated Ant-VEGF intra-vitreal injections</u>
b- Argon laser photo coagulation
c- Systemic steroid therapy


29 years old man




Retinitis Pigmentosa

The usual complaint   :

a- Day blindness
b- Color vision defects
<u>c- Nigh blindness</u>





27-years old man
C/O   lacrimation , irritation

Comment of the lids
What are the types and causes ?
What are the complications ?

Lid malpositions
Entropion and Trichiasis

- Trichiasis is rubbing of many lashes against the cornea; may occur alone or with entropion
- Entropion is inward rolling of lid margin
- Common causes are
  - Cicatricial
  - Involutional
  - Spastic
  - congenital
- Complications are mainly corneal (abrasions, ulcers and scars)
- Treatment is frequently surgical




Figure 3.88 Advanced pterygium

34-years old farmer
What is the pathogenesis ?
What are the indications for surgical removal ?
What are the commonest complications of surgery and how to minimize them?

it may recur , becomes more difficult and associates with symblepharon

Pterygium

- It is a degenerative and hyperplasic process in which a triangular fold of bulbar conjunctiva actively invades the cornea in the interpalpebral area
- Primary pterygium is that pterygium first time to appear in the patient eye.
- Pterygium activity depends upon the state of the advancing head
  - Active, Progressive = Head Serrated
  - Mild progressive = Head Rounded
  - Atrophic, inactive = Head is scared.



- Comment on conjunctiva & cornea
- What are the indications of surgical treatment ?

- **Answers:**

Diagnosis: Ptyregium

Comment :

Conjunctiva > conj. Epith. hyperplasia

Cornea >covered by apex of Ptyregium.

Indications of surgical ttt:

- If encroaches the pupillary area

- Progressive type

- Cosmetically annoying the patient



72-years old man




36-years old man

Comment of the lower eyelids

What is the underlying mechanism in both cases ?

Lid malpositions
Ectropion

- Rolling out of **lower** lid margin
- Unilateral or bilateral
- Mild (punctum eversion), moderate or severe
- Causes epiphora and chronic conjunctivitis
- May be
  - Congenital
  - Involutional
  - Spastic
  - Cicatricial
  - Mechanical
- Treatment is **mostly** surgical



4-years old female
A 6/36 OD , 2/60 OS

ame this sign
What are the management options for each eye? .

eatment is by

- Corneal cross-linking to strengthen corneal structur
- Eyeglasses and contact lenses in early stages
- Hard contact lenses
- Intracorneal ring segments
- Keratoplasty: DALK or PKP
- Refractive surgery is generally **contraindicated**

Keratoconus

- Is a disease condition in which the cornea bulges (**ectasia**) forwards acquiring the shape of a CONE due to weakness of corneal stroma
- It results in **progressive** myopic **astigmatism** that eventually becomes **irregular**
- It is diagnosed by
  - Cone shape of the cornea (MUNSON'S sign)
  - Scissors or spinning reflex in retinoscopy
  - Irregular placido disc image
  - Corneal topography showing marked steepening of the cornes >47 D with peak curve below the corneal center

- Treatment is by
  - Corneal cross-linking to strengthen corneal structure
  - Eyeglasses and contact lenses in early stages
  - Hard contact lenses
  - Intracorneal ring segments
  - Keratoplasty: DALK or PKP
  - Refractive surgery is generally **contraindicated**





Treatment of keratoconus











Figure 1.37 External hordeolum (sty)

Comment on the lids in both cases .
What is the pathology in each case ?
What is the management of each case?

SWELLINGS

- STYE:
  - acute suppurative inflammation of appendages related to cilia
  - points on lid margin
  - topical antibiotics and hot compresses
- ACUTE CHALAZION:
  - acute suppurative inflammation of a Meibomian gland
  - larger and more diffuse inflammation------→more **pain**
  - points on the conjunctival side
  - **oral** and topical antibiotics and hot compresses
- CHALAZION:
  - chronic inflammation of a Meibomian gland or follows an acute chalazion
  - localized with little edema
  - **incision and curettage from conjunctival side**


STYE
ACUTE CHALAZION
wiseGEEK


Chalazion
External Angular Dermoid

Chalazion / Meibomian cyst

- Sterile chronic granulomatous inflammation of a meibomian gland by retained sebum.
- Blepharitis is commonly present.





Chalazion:

Symptoms

- Subacute/chronic: gradually enlarging rounded nodule
- Acute chalazion: sterile inflammation or bacterial infection with localized cellulitis.


Signs

- A nodule within the tarsal plat
+/- inflammation.
- Points on conjunctival side



Chalazion:





Chalazion is better seen from the conjunctival side

Chalazion: treatment

1. Oral and topical antibiotics
2. Hot compresses several times daily
3. Vertical Incision and curettage---*video












Figure 3.35 Gelatinous superior limbal elevations with overlying fine white plaques (Trantas dots) in vernal limbitis


14-years old boy .
What is commonest presentation ?
What is the management of this case ?


Vernal keratoconjunctivitis

- Is a **recurrent** chronic conjunctivitis that aggravates in **spring** (spring catarrh) and summer
- It is very **common** in the middle-east and **Egypt**
- In hot months the eyes are red and **itchy** with a **ropy** discharge
- If lids are **everted** they show papillae that could be quite large and arranged as **cobble-stone** (palpebral VKH)
- The limbus may show **gelatinous** masses with white **Taranta** spots (limbal VKH)
- Corneal vascularization and ulcers may occur (**shield** ulcer)


Palpebral VKC with large papillae



Limbal VKC with Horner's spots


Severe pannus with corneal scarring


Shield ulcer
American Academy of Ophthalmology

- Treatment is preventive
  - Topical cromolyn sodium 4%
  - Prevents MAST cell degranulation
  - Should start 4 weeks before symptoms appear
- Then symptomatic in mild cases
  - Cold compresses
  - Dark glasses
  - Antihistamines/VCs
- Then topical Steroids in severe cases
- Steroids are dangerous with chronic use (more than 3x/day for 1 month)
- Topical immunomodulators as cyclosporine and tacrolimus may help spare steroids

IS this proliferative or not



A 50-year old female , diabetic for 17 years .


What is the recommended management ?

Diabetic retinopathy

- Affects many of the diabetics after an average of 5-7 yrs. in type II and 10-15 yrs. in type I
- A common cause of preventable blindness in the age of 30-65 years
- The retinopathy passes in distinct **stages**


1. **Non-proliferative** stages (simple or background retinopathy): mild□ moderate□ severe□ very severe (pre-proliferative)
2. **Proliferative** stage: formation of new vessels in response to retinal ischemia

Micro-angiopathy




Diabetic Retinopathy

Consequences of Chronic Leakage





Consequences of Retinal Ischemia





Manifestations


- **Microaneurysms**
  - the **earliest** manifestation
  - Dilatation of terminal capillaries due to loss of **pericytes**
  - Appear as red dots in colored photos and bright spots in FFA
- Retinal hemorrhages
  - Result from rupture of some capillaries
  - Are usually small and round (dot and blot)
  - Larger hemorrhages are in more advanced stage
  - Pre-retinal hge in proliferative
  - Appear red in colored photo and dark in FFA



Hemorrhages
Hard exudates
Microaneurysms



Cotton wool spots
Flame hemorrhage
Laser marks


Lipid
(Hard Exudate)
Cotton Wool Spot
(Soft Exudate)



Circinate
Exudate
(surrounds area of
swelling)


A
Fibrosis
Pre-retinal hge



NVD



00:44.93
Microaneurysms
NVEs
4/8/2008
30
E
a
c
b
d
Laser marks
NVDs
Laser marks


A
B
Retinal edema
C
D
E
F
G
H
I




Ischemia

- Hard exudates
  - Yellow deposits
  - May form rings (circinate rings)
  - Result from plasma lipids leaking from capillaries into retinal tissue
- Retinal edema
  - Focal thickening of retinal tissue
  - Indicate active leaking from vessels
- Macular edema
  - Edema and hard exudates within 1500 micron (1.5mm) from foveal center

- Cotton-wool spots
  - Fluffy white lesions
  - Result from minor vessel occlusion
  - Indicate ischemia
  - Sign of more advanced stages
- Venous changes
  - Dilatation
  - Looping
  - Beading
  - Signs of more advanced stages
- IRMA
  - Intra-retinal dilated capillary segments
  - Indicate pre-proliferative stage (severe and very severe)
  - Do not leak fluorescein


Preretinal hge

Fibrosis

- Capillary drop-outs
  - Areas of complete capillary loss in FFA
  - Indicate severe stage
- Neovascularization
  - New vessels budding from ischemic areas in response to Vascular endothelial growth factor (VEGF)
  - Can arise from disc surface (NVD) that bleed easily due to lack of vitreous support
  - Can arise from retinal surface (NVE) and result in recurrent hge or □
- Fibrous proliferation □
- Focal retinal tractional□
- Tractional retinal detachment
- Vitreous hemorrhage

Vitreous hge

Florid NVD




Treatment

- Management is a multidisciplinary approach with cooperation of the internist, endocrinologist and ophthalmologist
- Long-term control of blood sugar (HbA1c), serum lipids and ABP delay the appearance of retinopathy
- Treatment is stage-dependent and consists of a combination of
  - Medical treatment: multivitamins, antioxidants..etc
  - Laser treatment
  - Anti-VEGF
  - Vitrectomy

Laser treatment

- The commonly used technique is photocoagulation (heat-burning of retinal spots)
- Three main techniques
  - Focal treatment of micro-aneurysms
  - Macular grid for DME **Diabetic macular edema**
  - PRP (pan-retinal photocoagulation) for proliferative retinopathy **WITHOUT** traction retinal detachment
    - Rationale: photocoagulation of peripheral retina decreases secretion of VEGF and directs blood flow to central retina


360º PRP





Anti-VEGF


- Main drugs are bevacizumab (Avastin), ranibizumab (Lucentis) and aflibercept (Eylea)
- Block the action of VEGF produced by ischemic retina
- Used for
  - DME: repeated injections until edema resolves. Earlier treatment gives better results
  - Proliferative stage: before vitrectomy to reduce neovascularization
  - Neovascular glaucoma (NVG)


Vitrectomy

- Indications
  - Tractional retinal detachment when the macula is threatened
  - Unresolving vitreous hemorrhage
  - Bilateral vitreous hemorrhage
  - Vitreous hge in a patient who did not have PRP


Light source
Vitrector
Vitreous
Bleeding
Retina


a




b




56-years old female
What are the expected systemic associated diseases ?

Behçet's Disease


- Immune-mediated systemic vasculitis
- Presentation: 3^{rd}-4^{th} decade
- Most prevalent in Mediterranean region/Japan
- Associated with HLA-B51
- **Major diagnostic criteria**
  - Oral aphthous ulceration (100%)
  - Genital ulceration (80%)
  - Skin lesions (80%)
  - Uveitis (80%)
  - Other diagnostic criteria : arthropathy , thrombophlebitis






Retinal vasculitis


Hypopyon





61-years old male

What are the complications ?

C/BRVO

- Commonly occurs during sleep resulting in acute loss of vision in the morning
- Vision is usually 1/60-6/60
- Fundus is full of hemorrhages, cotton wool spots, tortuous looping veins and disc edema
- After the acute stage there is persistent macular edema
- After 3-4 months retinal neovascularization will occur and PRP **must** be done or neovascular glaucoma will occur
- Treatment is by monthly injection of anti-VEGF for 6-8 months


CRVO
Edward S. Harkness Eye Institute
Columbia University
BRVO


OCT showing macular edema
Fluid collecting within retina causing it to swell
Layers of Retina
A
Cotton wool spots
B
Nicking of V by A
Capillary drop outs



48-years old female
What are the emergency measurements?
What are the expected systemic causes ?

Retinal artery occlusion


- A 60-year-old had **sudden** darkening of vision in his left eye. He describes as if the room light has been suddenly shut down. He had a **previous** similar episode but lasted few seconds as he rubbed his eyes and the light came back. Now it is lost for 5 hours now. Vision is **HM** and the eye look very **normal**, however on closure of the right eye, the left pupil dilates. Examination reveals a generally **pale** retina and the fovea is standing out like a **cherry**-red spot. All retina arteries are **narrowed** and the blood column inside them seem segmented like **cattle**-trucks


Branch artery


- A 45-year-old noticed the appearance of a dark area in the upper field of his right eye. He is a drug-abuser. He is being treated for infective heart valve problems. His vision is 6/6 OU. Anterior segment free. Fundus examination shows an area of pallor along an attenuated inferotemporal arterial branch. Closer observation reveals an embolus lodged in the artery


C/BRAO


- Occlusion of a retinal artery is usually thrombotic in old age and embolic/vasculitic in young age
- Ischemia of all/part of retina results in ischemic necrosis of the inner ½ of retina (outer ½ receives) from choroidal diffusion
- Vision is lost (**HM**) if macula involved and positive scotoma if macula is spared
- Retina is whitish (cloudy swelling) esp. in posterior pole (several ganglion cell layers) and fovea stands out as CHERRY RED SPOT (no ganglion cells)
- Arteries are attenuated with segmentation of blood column
- Treatment (**EMERGENCY** 2-3 HRS) is mainly by acute **lowering** of IOP
  - IV acetazolamide and strong ocular **massage**
  - Paracentesis
  - Antiplatelets/fibrinolytics
  - VD by breathing 5% CO_2


Milky white central retina
Cherry red fovea


Cattle trucking of vessel
Cherry red spot


Ischemic retina


BRAO



32 years old female

What is the expected refraction ?

What are the expected complications ?



42 years old man , working on drilling machine

Past history of trauma to the left eye with and FB hitting the eye

Describe the findings


What is the possible underlying cause ?

Blunt trauma

- Posterior segment manifestations
  - Vitreous: trauma is a common cause of vitreous hemorrhage
  - Retina
    - Retinal hemorrhage
    - Terson syndrome
    - Purtscher’s retinopathy
    - Commotio-retinae (Berlin’s edema)
    - Retinal tear
    - Retinal Dialysis
  - Choroid
    - Choroidal rupture
  - Optic nerve
    - Traumatic optic neuropathy
    - Optic nerve avulsion

Intraocular foreign body (IOFB)

- Iron: Occurs in manual workers who use hammers without protective goggles
  - A small fragment penetrates the eye causing physical injury and later chemical injury (siderosis bulbi)
  - Infection is rare (heat generation)
  - X-ray is done or ultrasound. CT scan is also useful
  - **MRI is CONTRAINDICATED if IOFB is SUSPECTED**
  - **FB** is removed by vitrectomy and FB
  - **Siderosis** occurs months later with rusty discoloration of the iris and lens, secondary glaucoma, night blindness, retinal pigmentary changes and low ERG
- Copper (**chalcosis**); acute stage picture of sterile endophthalmitis, later blue-green stain of ocular tissues
- Glass: inert, small pieces can be left inside the eye if hard to remove


FB
Ocular siderosis
DFOV 15.1cm
STND/I
512
MF:1.2
L
R
kV 120
mA 150
Head



1. This is musca volitantes
2. This lesion could result from retinal detachment




30. The acute red eye shows signs of angle closure glaucoma
31. This patient needs topical atropine



32.There are features of fungal corneal ulcer T

33.Topical acyclovir is the best treatment



VKC

Severe pannus with corneal scarrin


38.The everted lid is likely to show conjunctival follicles
39.Itching and photophobia are the main symptoms

Trachoma

- Is a chronic conjunctivitis caused by ***Chlamydia trachomatis*** *serotypes A, B and C,* endemic in many parts of the world including **Egypt**
- The organism is acquired during **childhood** by flies or dirty hands, many cases can be prevented by healthy habits as hand and face washing
- Conjunctiva shows **both** follicles (lymphocytic aggregates) and papillae (epithelial proliferation)
- Cornea may be involved with **follicles** (Herbert's rosettes) and superficial vascularization (**Pannus**) in the upper cornea

WHO classification


- The WHO grading system for trachoma classifies the disease in 5 grades:
  - Trachomatous Inflammation: Follicular (TF) - which mostly requires topical treatment.
  - Trachomatous Inflammation: Intense (TI) - during which topical and systemic treatments are considered.
  - Trachomatous Scarring (TS): when scars are visible in the tarsal conjunctiva and may obscure tarsal blood vessels.
  - Trachomatous Trichiasis (TT): when an individual is referred for eyelid surgery.
  - Corneal Opacity: a stage during which a person is irreversibly blind.


Halbertstaeder Prowazek eosinophilic inclusions
Elementary body
PTCs AND PTDs
Arlt’s line
Subconjunctival fibrosis
Shallow fornix
Posterior symblepharon
EyeRounds.org
Trichiasis

Follicles

Follicles and papillae


a




b




c




d



e




f



Arlt’s line (fibrosis)

Corneal scarring
Trichiasis

Pannus siccus


PANNUS ANNULOSUS
Scar
Cellular infiltration
Superficial
corneal
vascularization


- Infection remains for many years resulting in **progressive** fibrosis manifested as
  - Conjunctival fibrosis
  - Shallow fornix
  - Symblepharon
  - Trichiasis
  - Entropion
  - Corneal ulcers and scarring
- Treatment is by
  - Oral tetracycline for 2-3 weeks to eradicate infection
  - Oral erythromycin in children
  - Azithromycin single dose of 1gm in adults and 20mg/kg in children
  - Treatment of complications esp. trichiasis and entropion

TRACHOMA GRADING CARD

– Each eye must be examined and assessed separately.
– Use binocular loupes (x 2.5) and adequate lighting (either daylight or a torch).
– Signs must be clearly seen in order to be considered present.

The eyelids and cornea are observed first for inturned eyelashes and any corneal opacity. The upper eyelid is then turned over (everted) to examine the conjunctiva over the stiffer part of the upper lid (tarsal conjunctiva).


The normal conjunctiva is pink, smooth, thin and transparent. Over the whole area of the tarsal conjunctiva there are normally large deep-lying blood vessels that run vertically.




Normal tarsal conjunctiva (x 2 magnification). The dotted line shows the area to be examined.

TRACHOMATOUS INFLAMMATION – FOLLICULAR (TF)*: the presence of five or more follicles in the upper tarsal conjunctiva.*

Follicles are round swellings that are paler than the surrounding conjunctiva, appearing white, grey or yellow. Follicles must be at least 0.5mm in diameter, i.e., at least as large as the dots shown below, to be considered.






Trachomatous inflammation – follicular (TF).

TRACHOMATOUS INFLAMMATION – INTENSE (TI)*: pronounced inflammatory thickening of the tarsal conjunctiva that obscures more than half of the normal deep tarsal vessels.*


The tarsal conjunctiva appears red, rough and thickened. There are usually numerous follicles, which may be partially or totally covered by the thickened conjunctiva.



Trachomatous inflammation – follicular and intense (TF + TI).

TRACHOMATOUS SCARRING (TS)*: the presence of scarring in the tarsal conjunctiva.*


Scars are easily visible as white lines, bands, or sheets in the tarsal conjunctiva. They are glistening and fibrous in appearance. Scarring, especially diffuse fibrosis, may obscure the tarsal blood vessels.



Trachomatous scarring (TS)

TRACHOMATOUS TRICHIASIS (TT)*: at least one eyelash rubs on the eyeball.*


Evidence of recent removal of inturned eyelashes should also be graded as trichiasis.



Trachomatous trichiasis (TT)

CORNEAL OPACITY (CO)*: easily visible corneal opacity over the pupil.*

The pupil margin is blurred viewed through the opacity. Such corneal opacities cause significant visual impairment (less than 6/18 or 0.3 vision), and therefore visual acuity should be measured if possible.



Corneal opacity (CO)

TF:– give topical treatment (e.g. tetracycline 1%).
TI:– give topical and consider systemic treatment.
TT:– refer for eyelid surgery.

WORLD HEALTH ORGANIZATION
PREVENTION OF BLINDNESS AND DEAFNESS

Support from the partners of the WHO Alliance for the Global Elimination of Trachoma is acknowledged.





54.MRI is contraindicated
55.Vitrectomy is essential




56. Treatment is by phakoemulsification
57.Biometry is essential

Stage III




66. This could be grade II papilledema

67. If papilledema vision should be poor



68 this patient could have a parietal lobe mass. temporal

69.The lesion is on the right side of the brain



- What is the diagnosis?
- Complications ?

- **Answers**

Diagnosis :

Entropion of lower lid with trichiasis

Complications:

Chronic conjunctivitis,

conjunctival scar, corneal ulcer &



- What is the diagnosis?
- Treatment ?


- **Answers**

Diagnosis :

Cicatricial ectropion

Treatment :

V to Y plasty or Z plasty



- - What is the diagnosis?

- Surgical operation in the lid ?

- **Answers**
- **Entropion in lower lid, trichiasis & corneal ulcer**
- **Surgical operation:**
  - **Lateral canthotomy, lateral canthoplasty**



- Diagnosis
- treatment

- **Answers :**

-Stye

ttt of P.F. “ staph. Aureus”

-local antibiotics & eye drops

-Hot fomentation



- Diagnosis
- Complications
- treatment
- **Answers:**

-Diagnosis: Ophthalmia neonatorum

-Complications:

Corneal ulceration , iridocyclitis

-ttt:

-Prophylactic ttt.

-Curative ttt: lotions, topical and systemic antibiotics and atropine ointment in




- Diagnosis
- etiology

- **Answer:**

Phlyctenular conjunctivitis (limbal phlycten)

Etiology :

Hypersensitivity reaction to endogenous antigens e.g. bacterial antigens as T.B & chlamydia.



- Diagnosis
- treatment

- **Answers:**

-Bulbar spring catarrhal

ttt :topical steroids, mast cell stabilizers,

anti histaminic

Dark glasses & cold compresses.



- Diagnosis
- 2 causes

- **Answers:**

-phlycten

-Causes:

Hyper sensitivity to an endogenous antigen e.g. tuberculo-protein, Intestinal parasites, staphylococcal blepharoconjunctivitis.



- Comment on lashes
- complications

- **Answers:**

-Ulcerative blepharitis

-Complications:

Chronic conjunctivitis, Madarosis,

trichiasis, ptylosis, epiphora,

Ectropion, corneal ulcer.

-



- What is this sign called?
  - pathogenesis

- **Answers:**

Sign:

Marcus –Gunn phenomenon

Pathogenesis:

Faulty Innervation

“motor fibers from 5th nerve reach levator instead of the 3rd nerve”



- Diagnosis
- treatment

- **Answers:**

Diagnosis:

mucopurulent conjunctivitis

Ttt:

-Eye lotions

-Antibiotics ointments e.g. tobramycin at night

-Antibiotic eye drops

-Hot foments



- Cause of this appearance
- 2 eye drops

- **Answers:**


Cause:

post. Synechiae

2 eye drops:

Atropine sulfate

corticosteroids



- Diagnosis
- Name 2 causes


- **Answer**

Diagnosis :

rubeosis iridis – Peripheral iridectomy

Causes :

Diabetic Retinopathy and CRVO



- Diagnosis
- What is the visual complaint?

- **Answers:**

Diagnosis:

Irido-dialysis

Visual complaint :

Uniocular diplopia



Diagnosis


- Factors affecting prognosis
- **Answers:**

Diagnosis:

Lt. congenital ptosis

Factors affecting prognosis:

- **Amount of ptosis**
- **Extent of levator function**
- **If 3^{rd} nerve palsy >> correct squint first**
- **If 5^{th} nerve palsy >> postpone the op. till 5^{th}**




- A 68y patient complaining of sudden diminution of vision.
- What is the Diagnosis?
- mention two systemic predisposing condition

- **Answers:**

Diagnosis: CRVO

2 Systemic P.F.:

Systemic hypertension & Diabetes mellitus



- Diagnosis
- Antiviral drugs for ttt


- **Answers:**

Diagnosis:

Herpetic corneal ulcer “Dendritic ulcer”

Antiviral drugs:

Acyclovir , vidarabine , T3F & IDU



- Diagnosis
- What is the suspected refraction of this patient?
- **Answers:**

Diagnosis:

Keratoconus " Munson`s sign"

Suspected refraction:

axial myopia & Astigmatism.




- Diagnosis
- 2 posterior segment diseases cause it

- **Answer**

Diagnosis: rubeosis iridis

Causes: Diabetic Retinopathy and CRVO



- Diagnosis
- Deferential diagnosis

- **Answers:**

Diagnosis : **CRAO**

D.D.: **"For cherry red spot"**

- **Commotio retinae**
- **Quinine poisoning**
- **Macular hole surr. By RD**
- **Amauratic family idiocy**



- Diagnosis
- Expected field of vision
- **Answers:**

Diagnosis:

Glaucomatous cupping of optic nerve

- Expected field of vision:
- Tubular field.



- Diagnosis
- 2 syndromes associated with it
- mention effect on optic nerve

- **Answers:**

Retinitis Pigmentosa

2 Syndromes :

Bardet - biedl syndrome

Refsum's disease

Effect on optic n.:

Waxy disc pallor due to consecutive optic atrophy



- The optic disk of this patient show…….
- Name a cause for this condition

- **Answers**

Comment :

Papilleodema

Cause:

Elevated intracranial tension.



- A 57y patient with sudden painful drop of vision IOP is stony hard
- Name 2 medication for emergency
- treatment of this case


Answer

- **Diagnosis: Acute congestive glaucoma**
- **2 Medications for emergency : hyper-osmotic agent, topical miotics, topical steroids**
- **Ttt: ttt essentially surgical recent….surgical iridectomy late….their is PAS ,an external fistulizing operation .**



- Nerve& muscle affected ?
- Direction of gaze which diagnose this case ?
- The main complaint of the patient


- **Answer**
- **Nerve& muscle affected: Rt Abducent nerve-RT. Lateral rectus**
- **Direction of gaze: To the right**
- **Main complaint: Binocular Diplopia**



- Diagnosis
- Component of it

- **Answers:**

Diagnosis:
hypermetropia
Components:
Total , Latent , Manifest
hyperopia



- Type of squint & its angle
  - Confirmatory test

- **Answers:**

Type of squint :

exotropia

angle:

30

Confirmatory test:

Cover test



- Diagnosis
- treatment

- **Answers:**

Diagnosis:

After cataract

TTT:

- No interference if vision is not affected

- If thick : surgical intervention

- Diagnosis
- treatment



- **Answers:**

Diagnosis:


myopia

TTT:

- eye glasses with concave minus lenses

- contact lenses

- refractive surgery if indicated



- What s this inustrument called
- Used for........


- **Answers:**

Instrument:

Applanation tonometry (gold mann)

usage:

IOP measurement



- What is the upper lid abnormality?
- What are the complications?


- **Answers:**

Comment :

Left upper lid ptosis

Complications:

Amblyopia & Squint scoliosis and ocular torticollis.



- Comment on lens
- This is an association of………..syndrome


- **Answer**

Comment :

lens subluxation

Syndrome:

Marfan’s syndrome



- A 68y old woman with cataract extraction . Complaining of drop of vision which was managed
- What was the cause of drop of vision?
- What was the management?

- **Answer**

Cause:

posterior capsular opacification (after cataract)

Management:

YAG laser capsulotomy



- Comment on the lower lid
- Name 2 possible complications of this conditions

- **Answers:**


Comment:

Senile ectropion

2 possible complications:

xerosis

corneal ulcer



- Diagnosis
- Name 2 surgical procedures for ttt of this condition

- **Answers:**


Diagnosis :

Buphthalmos

2 Surgical procedures:

-goniotomy

-trabeculotomy



- What are these field defect called?
- Name the cause

- **Answer**


Field defect:

Bitemporal hemianopia

Cause:

Optic chiasma lesions (nasal fibers damage)

e.g. Pituitary gland



- What's the error of refraction in this patient ?
- What're the complications of this case ?

- **Answer**

Error of refraction :

High myopia

Complications:

Chorio-retinal degenerations

retinal tears

retinal detachment



- Diagnosis
- 3 causes

- **Answers**

Diagnosis:

Symblepharon

Causes:

- Post-trachomatous

- Post-operative (Pterygium excision)

- Ocular cicatricial pemphigoid



- The eye lid & Conjunctiva show
    - possible findings in the crystalline lens


- **Answers**

Comment :

Ecchymosis & subconjunctival hemorrhage"

Possible findings:

(Concussion cataract " Rosette-shaped" - Lens subluxation or dislocation)



- Diagnosis
- 2 causes

- **Answers:**

Diagnosis :

Lens subluxation

Causes:

Marfan's syndrome

Homocystenuria &
Trauma



- What's the sign called ?
- 3 possible causes

- **Answers**

Sign :


Leukocoria

Causes :

Retinoblastoma

congenital cataract

Retinpathy of prematurity



- What is the error of refraction ?
- How to correct ?


- **Answers**

Error of refraction:

Hypermetropia

Correction :

Spherical Convex "plus" Lens



- Diagnosis
- **mention caustive organism**

- **Answers:**

Diagnosis :

Hypopyon corneal ulcer

Causative organism:

pneumococci



- Diagnosis
- Mention disease cause this


- **Answers:**

Diagnosis:

exophthalmos

Disease :

Hyperthyrodism



- Diagnosis
- treatment

- **Answers:**

Diagnosis: Corneal foreign body

TTT: Surgical removal



- Diagnosis
- **mention muscle affected and its nerve supply**

- **Answers:**

Diagnosis: **Left upper lid ptosis**

Muscle affected :

levator palpebrae sup.

innervation:


oculomotor n.



- Diagnosis
- Mention 2 complications

- **Answers:**

Diagnosis:
sublaxated &cataractous lens
Complications:
lens dislocation
2ry Glaucoma
Iridocyclitis



- Diagnosis
- Mention 2 complications

- **Answers:**

Diagnosis: Blood staining of the cornea “total hyphema or 8-ball hyphema”

2Complications:

Elevation of IOP

Corneal staining



- Diagnosis
- Mention 2 ttt

- **Answers:**

keartic precipitates

ttt :

Topical : Atropine sulfate & corticosteroids.

Systemic: systemic steroids (in severe cases)


& Antibiotics (in infective cases)

- **sederosis bulbi**
- **Patient with foreign body in his eye from one year**
- **In picture you will see one eye normal &other eye(black iris)**

A case of recent open globe injury:

- Primary care would be sterile dressing and prophylactic antibiotic eye drops F
  Only patching


- 1ry repair as soon as possible is mandatory. T



Iris prolapse

A patient with blunt ocular trauma:


- Diplopia is a possible manifestation.
  T
- This is the only ocular sign can be seen. F



Iridodialysis

This case of complicated cataract:


- Commonly caused by systemic disease. F
- Can be associated with retinal detachment. T



Traumatic cataract (flower shaped)

- *The first line of treatment in acid burns of the eye is Instilling a drop of surface anesthetic into the eye*


- *false*



Chemical acid burn

Male patient 30 years old complains of recurrent attacks of obscuration of vision and headache his fundus examination showed this picture.

- The most important investigation is FA. ايه بقى False
- In early stages of this disease vision may be 6/6. T



papilledema

A patient with recent rhegmatogenous retinal detachment:

- **fluid accumulates between Retinal pigment epithelium and Bruch’s membrane. T**
- Hypermetropia is a predisposing factor. F



Rhegmatogenous RD


In this case of corneal ulcer:

- **Irrespective of the aetiology of a corneal ulcer, Corticosteroid eye drops are contraindicated.**

T

- Corneal ulceration is caused by injury to the third cranial nerve.

F



Fundus picture of 30 years old female:

- **most common presentation is diplopia.** ايه بقى **F**
- Decrease of vision caused by dystrophies of the retinal rodes.

T



Ritinitis pegmentosa


Field change of 50 years old patient

- Examination of IOP is important point of in diagnosing this case.

  T

- In early stages of this disease cup/disc is vertically oval.

  T




Ten years old boy complains of itching. On examination, there are with large flat topped cobble stone papillae raised areas in the palpebral conjunctiva is:

- Most common cause is trachoma.

  F

- **Ptosis is a possible manifestation of the disease.**

  T



Vernal keratoconjunctivitis

A case of post trachomatous degeneration.

- Common causative organism is *Chlamydia trachomatis serotypes D to K.*

  F

- *The patient is infectious in this stage.*

  F



Tarsal conjunctive.

- In this stage there is active inflammation of the disease.

F

- This lesion may obscure tarsal blood vessels.

T



Trachoma (scarring phase)

This is an eye of 30 years old male:


- Conservative treatment is recommended in this case.

F

- It may be caused by exposure to infra- red rays

UV rays

F



Ptregium


55 years old patient with unilateral sudden loss of vision. He is diabetic since 2 years, ocular examination revealed this picture, fundus of the other eye is within normal limits.

- Likely diagnosis is proliferative diabetic retinopathy.

  F

- There is high risk of developing glaucoma within next 3 months.

  T



CRVO

A young patient with sudden painless loss of vision, with systolic murmur and ocular examination reveals a cherry red spot with clear AC, :

- **the likely diagnosis is Branch Retinal Vien Occlusion.**

  F

- **cherry red spot is due to increased choroidal perfusion.**

  F



CRAO

Case 1:



A 12-years-old boy presented with red aching eye. **What is the most likely underlying pathogenesis of this disease?**

a. Type 1 hypersensitivity.
b. Viral infection.
c. Type 4 hypersensitivity.
d. Trauma.
e. Hypoxia (Contact lens wear).

Case 2:



A 28-years-old man presented with painful, swelling in the upper lid . He gave a history of repeated similar condition in both eyes over the past 6 months. **Which investigation is most relevant to this patient?**

a. Bleeding and coagualtion profile.
b. Blood sugar levels and Hb A1c.
c. Blood test for anemia and Hb.
d. Tuberculin skin reaction.
e. Serum IgE.

Case 3:




Acute dacryocystitis?

A 48-year-old woman with presented with pain , epiphora and discharge . **What is the best management?**

a. Topical antibiotic drops.
b. Warm compresses.
c. Incision and drainage.
d. Topical steroids.
e. Systemic steroids.


Case 4:



A 38-years old man , presented with red painful eye and drop of vision. **What is most correct about the prognosis?**

a. Patching produces symptomatic relief and cure.
b. It has a self-limiting course.
c. Corneal perforation is likely if untreated.
d. Corneal hypoesthesia will occur.
e. The cornea can clear after successful treatment

Case 5:



مش دي صورة phlycten?

A 48-years-old woman presented with acute red eye . She complains of pain and swelling in shoulders and knees. **What is the most commonly associated eye disease with this condition?**

a. Keratoconjunctivitis sicca.
b. Central corneal ulcer.
c. Acute anterior uveitis.
d. Corneal pannus.
e. Retinal detachment.

Case 1:



Which is the correct pair?

a. Corneal sensation, Esthesiometer.
b. Tear film stability, tear break-up time.
c. Degree of levator action, Burke’s (ruler) test.
d. Dry eye, Schirmer’s test.
e. Esotropia, Hirschberg test.


Case 2:



What is the pathology?

a. Corneal scarring.
b. Calcium deposition.
c. Conjunctival invasion of cornea.
d. Corneal infiltration.
e. Corneal vascularization.

Case3 :




rhegma

48-year-old myopic male patient, complaining of flashes and floaters a week ago followed by curtain in the lower field of vision . What is the next step in management?

a. Bed rest and Observation
b. Urgent surgery
c. Systemic steroid therapy
d. Perimetry.
e. Fundus Fluorescein angiography.

Case 4 :





This is the fundus picture and lips of an 28-years old patient , presenting with acute drop of vision. **Which is the least characteristic feature of this disease?**

a. HLA typing help to reach the diagnosis.

b. The condition is more common in males .

c. The eye involvement may be the first presentation.

d. The prognosis for vision is good.

e. Steroids are the mainstay of treatment

Case 5 :



2-year old boy . Parents noticed change in appearance.

What is the next step in management?

RB?

a. Fundus Fluorescein Angiography

b. Antibiotic eye drops.

c. Ocular ultrasound.

d. Cataract surgery .

e. Urgent glaucoma surgery .


Case 1 :



72-years old man presented with a complaint " my eyelids gets on my way as the day passes". He also reports attacks of horizontal diplopia in the evening. **What is the next step in the management?**

a. Fasting blood sugar and Hb A1c.
b. Refer for levator muscle strengthening procedures.
c. EMG.
d. Chest X ray.
e. Steroid therapy.

Case 2:



ليه مش papilledema?

A 60-year old male has left **acute** drop of vision to 1/60 and loss of color vision. **What is the most likely diagnosis?**

a. Anterior ischemic optic neuropathy


b. Acute papilledema.

c. Acute glaucoma.

d. Choroidal neovascular membrane.

e. Toxic optic neuropathy

Case 3:



Optic nerve head of a 56-years-old man. He presented to the outpatient clinic for routine check up. Pupillary light reflexes were normal in both eyes. **What is the next choice?**

a. Fundus Fluorescein Angiography.
b. Visual field testing.
c. Observation and repeated fundus examination .
d. Systemic steroid therapy.
e. Systemic diuretics as Lasix.

Case 4:




What is this instrument used for?

a. Corneal curvature estimation.
b. Palpebral fissure width estimation.
c. Corneal sensation measurement.
d. Exophthalmos measurement.
e. Intra-ocular pressure measurement.

Case 5 :

Diabetic maculopathy?



A 39 years old female, diabetic for 8 years , Her last HB A1c is 8.5% . BCVA 6/24. **What is your management choice for this case?**


Hard exudates?

a. Pan-retinal photocoagulation
b. Close follow-up.
c. Intra-vitreal injection of Anti VEGF.
d. Pars-plana vitrectomy.
e. Strict diabetic control.

Clinical station 4

Trauma

Case 1 :



This 17 -years-old boy presented with double vision. He denies history of trauma to this eye. **What is your logical choice for this patient?**

a. Refraction and eye-glasses.

b. Cardiologic consultation.

c. Corneal topography and LASIK.

d. Topical steroids.

e. Neurologic consultation.

Case 2:


Case 3 :



A 23-years-old man presented to the ER with a history of severe blunt trauma to the eye. **What is the best management choice?**

a. Patching the eye with topical antibiotic eye drops and ointments
b. Systemic antibiotics
c. Topical anesthesia to relief the pain
d. Emergency surgery
e. Topical cycloplegia

Case 3 :



A 20-years old with history of right eye **trauma** by tennis ball. He notices flashes and black spots in his field of vision. This is his fundus examination. **What is your next step in management?**

a. Reassurance and follow-up in two weeks
b. Systemic steroids
c. Argon laser application (Laser retinopexy).
d. Systemic antibiotic
e. Panretinal photocoagulation

Case 5 :



What is the most serious complication of this condition if left untreated?


a. Neovascular glaucoma.

b. Phacomorphic glaucoma.


c. Iridocyclitis.

d. Lens dislocation.

e. Corneal blood staining.



1- This is the pathway for **vitreous** circulation.
2- The fluid is secreted by the **iris** processes.
3- the uveoscleral outflow is the **main** route for drainage.
4- The fluid drains into the angle of the **posterior** chamber.




5- This chart measures one property of vision.
6- It can be used to assess visual acuity.
7- Chart can be used to assess field of vision.
8- There is a colored version of the chart.



9- These are the openings of **Zeis** glands.
10- This photo shows a case of **blepharitis**.
11- This condition has **corneal** complications.
12- This condition is one of the causes of **dry** eye.

An 80-year-old with fair skin




13- This ulcer is **malignant**.

14- It is more common in the **lower** lid.

<u>15- Main line of treatment is **radiotherapy**. F</u> surgery


<u>16- Lymph node **metastasis** is early. F</u> direct invasion

A 2-year-old with acute bilateral lid hemorrhage



17-This is a **primary** orbital disease.
18- **Abdominal CT** is an important investigation.
19- **This** is a **bleeding** disease not a neoplastic disease.
20- **The** arrows point to a **retinoblastoma**.

A 10-year-old with gradual loss of vision



21- The optic nerve is pale. T

22- There is **left** orbital mass.


23- **The** mass lies in the muscle **cone**.

24- The mass is of **glial** cell origin.

A 40-year-old on hydroxychloroquine



29- This lesion is described as **diffuse**.
30- **T**his is nodular **conjunctivitis**.
31- Treatment is by topical **antibiotics**.
32- This patient commonly has **collagen** disease.



33- These are conjunctival **papillae**.
34- **Treatment** is by oral **azithromycin**.
35- **Trichiasis** is a known complication.
36- The condition is **secondary** to **Chlamydial** infection.

Field of vision of a 60-year-old



37- This patient has an **optic** nerve disease.
38- This defect is called **hemianopia**.
39- This lesion could result from **cerebral vascular** occlusion.
40- The other eye has a similar defect.




41- This is **apparent** squint.
42- This is **exophoria**.
43- The angle of squint is **30º**.
44- Treatment is by **convex** glasses.



45- There is right **esotropia**.
46- This baby will complain of **diplopia**.
47- Hirschberg test shows **no** squint.
48- The is a **telecanthus**.

A 50-year-old with gradual loss of vision



49- The cup/disc **ratio** is 0.8
50- There is **notching** of the neural rim.
51- The lamina cribrosa is **visible**.
52- This patient is **glaucomatous**.

A 70-year-old with gradual loss of vision



FHM


53- This patient can see maximum 6/60.

54- **Ultrasonography** is likely needed.

55- **Biometry** is a likely investigation.

56- **Amsler** grid is useful in diagnosis and follow up.

A 40-year-old with acute drop of vision




57- This patient is likely **anemic** .

58- This patient is likely **thrombocytopenic**.

59- This patient could be **leukemic**.

60- Arrows point to a **subhyaloid** hemorrhage.

A 70-year-old with sudden loss of vision



61- **MRI** may show **bilateral** enlargement of the medial and inferior recti.
62- The optic disc is **pale**.
63- Field may show an **altitudinal** defect.
64- Emergency **IV steroids** are needed.

cong. glaucoma



- This baby presented with right watery eye and photophobia. What is your first step to establish the diagnosis?

a) Syringing of the lacrimal system F

b) IOP measurement under GA (+ corneal diameter) ^T

c) TTT….. Surgical T

Retinoblastoma



- This a fundus picture e of a 2-year old child

a) Intravitreal antiVEGF is the best treatment option F   chemo +radio

b) The lesion is caused by intrauterine infection. F


Developmental cataract RB??

- What is true regarding this child who presented with white pupil?

a) Congenital glaucoma is the commonest cause. F

b) B-scan is the most useful investigation to be done. T ???





Developmental cataract

- What is the best timing for removal of this lens opacity?

a) At 4 years when the eye size is adequate. F

b) We can wait 6 months as it is unilateral. F


c) As early as possible.T

Keratoconus
Munson sign



- 21- year old presented with drop of vision with frequent change of eyeglasses, which of the following is true?

a) The condition is caused by weakening of the zonule F
b) This sign shown is called oil droplet sign.F
c) The condition is usually bilateral.T
d) With repeated change of eye glasses. T



Corneal tomography before restrictive surgery


a) This investigation is optical coherence tomography. F

b) It is usually performed before cataract surgery. F

c) It is the benchmark in corneal ectasia diagnosis and follow up. T

- before cataract surgery___ biometry . **A scan> keratometry**

chorioretinal



-With pigmentation – pathological myopia
-These may be the result of an old infection

22-year old patient has this fundus picture ALL FALSE


a) This lesion could be due trauma.
b) Expected refraction of such patient is -2
c) The patient has poor unaided VA but corrected VA could be 6/6



- The mother of this child came complaining about his look. What would you tell her?

a) He has left esophoria. F

b) Epicanthus is the cause of this abnormal look. F



- This is the picture of a five year old before and after wearing his glasses. Which of the following is true?


a) He needs left medial rectus recession. f

b) The expected refraction is -5.00. F Hypermetropia



- What does this ocular motility examination most likely suggest?
- Left concomitant esotropia
- Right esophoria
- Right paralytic esotropia

Left Homonomys hemianopia with macular sparing



1. Dual blood supply is responsible for this phenomenon (T)
2. Lesion is due to affection of parietal lobe (F)
occipital

Ischemic optic neuropathy






3.Patient complains of defective vision (T)

4. In adults the most common cause is infectious (F)

- In giant cell arteritis (atherosclerosis- thrombosis)

papilledema




5. Lumbar puncture opening pressure can be 45cm H2O for this patient (T)

6. Pupillary reaction is normal (T)

- the most dangerous cause is intracranial tumor

Rt eye esotropia(6th CN palsy)



7. Patient complains of lagophthalmos (F) in 7th cn palsy

8. Diplopia increases on looking upwards (F) on looking right

-MRI indicated

-IOP monitoring ???

chalazion



9. Treatment is essentially conservative (F) Surgical ttt

10. Causative organism is demodex folliculorum (F) staph

- demodex folliculorum cause parasitic blepharitis

Congenital ptosis



11. Treatment is postponed till school age (F) ???

12. Condition is essentially unilateral (F) ???

Congenital ptosis is the most common type of ptosis

-ttt if mild__ follow up
severe___ surgery(frontalis sling)

-amblyopia is a complication


Lower lid entropion



13. Vision is affected if treatment is neglected (T) by corneal ulcer
14. Tarsorrhaphy is the treatment of choice (F) Weiss operation
- Tarsorrhaphy is the surgery of 7th cn palsy

Tear drop sign

Blow out fracture



15. The IR is the entrapped muscle (T)
16. The diplopia increases on looking downwards (F) upwards ???

- Vertical diplopia

MRI unilateral Proptosis



17. Lesion is intraconal (T)
18. Non axial proptosis is the main presentation (F)

Bilateral proptosis
Thyroid eye ds



19. Thyroid function test is needed (T)

20. Patient may complain of color vision affection (T)